

The history and purview of phylogeography: a personal reflection

JOHN C. AVISE

Department of Genetics, University of Georgia, Athens, GA, 30602–7223, USA

Abstract

Last year marked the 10th anniversary of the birth of phylogeography as a formal discipline. However, the field's gestation began in the mid-1970s with the introduction of mitochondrial (mt) DNA analyses to population genetics, and to the profound shift toward genealogical thought at the intraspecific level (now formalized as coalescent theory) that these methods prompted. This paper traces the early history and explosive growth of phylogeography, and closes with predictions about future challenges for the field that centre on several facets of genealogical concordance.

Keywords: allelic pathways, coalescent theory, gene trees, genealogical concordance, intraspecific phylogeny, mitochondrial DNA

Introduction

Phylogeography is a field of study concerned with the principles and processes governing the geographical distributions of genealogical lineages, especially those at the intraspecific level. The word itself was coined a decade ago (Avice *et al.* 1987a) and its use in the evolutionary genetics literature has grown exponentially since then (Fig. 1). As of the end of 1996, more than 130 papers had employed 'phylogeography' in the title or as an index word, and they represent only the tip of the iceberg because numerous additional studies have dealt with the topic implicitly although not by name. As a subdiscipline of biogeography (Fig. 2), phylogeography emphasizes historical aspects of the contemporary spatial distributions of gene lineages (Avice 1996a). The analysis and interpretation of lineage distributions usually requires input from molecular genetics, population genetics, phylogenetics, demography, ethology, and historical geography. Thus, phylogeography is an integrative discipline.

In purest form, empirical phylogeographic analyses deal with the spatial distributions within and among populations of alleles whose phylogenetic relationships are deduced. Because mitochondrial (mt) DNA evolves rapidly in populations of higher animals and usually is transmitted maternally without intermolecular recombination, it has been the workhorse of most (> 80%) of the

phylogeographic studies conducted to date (Fig. 3). However, empirical or theoretical treatments that address phylogenetic aspects of the spatial distributions of any genetic traits (morphological, behavioural, or any other) also can qualify as phylogeographic under a broader definition of the term. Furthermore, a matrilineal phylogeny (or any other allelic transmission pathway) constitutes only a minuscule fraction of the composite genealogical information within a sexual pedigree (Fig. 4). A phylogeny for spatially structured populations can be conceptualized as a statistical distribution of partially bundled allelic pathways of descent (Fig. 4) each characterized by its own unique coalescent pattern (Maddison 1995; Avice & Wollenberg 1997). The many distinctions yet connections between notions of phylogeny at the levels of genes vs. populations have made phylogeography a rich point of contact between the traditionally distinct fields of population genetics and phylogenetic biology (Avice 1989a; Hey 1994).

To introduce this special issue of *Molecular Ecology* on phylogeography, I will recount briefly the history of the discipline from a personal, anecdotal (and no doubt biased) perspective. Phylogeographic efforts have been tied closely to analyses of animal mtDNA, so that is where the story will begin.

History of phylogeography: one researcher's view

Science often is serendipitous, as the following stories well illustrate. Shortly after joining the University of Georgia as an Assistant Professor in 1975, I gave a departmental

Correspondence: J. C. Avice, Tel.: +01-706-542-1456; Fax: +01-706-542-3910; E-mail: Avice@bscr.uga.edu




Fig. 1 Published articles with 'phylogeography' or 'phylogeographic' in the title, or as index terms, following the introduction of these words in 1987. The number of such papers has approximately doubled across each successive 2-year time interval. This computer search was conducted in October 1996, and included listings in Current Contents, Biosis, and the Expanded Academic Index.

biogeography

ecogeography	phylogeography	
e.g., Bergmann's rule	Dispersal	Vicariance
(tends to emphasize the influence of contemporary natural selection on geographic distributions of traits)	(tends to emphasize the influence of historical factors on geographic distributions of gene lineages)	

Fig. 2 The heuristic place of 'phylogeography' within the framework of biogeography (after Avise 1994). Of course, ecogeographic and phylogeographic perspectives are not mutually exclusive because natural selection is among the historical factors that also influences lineage distributions.

seminar describing work on allozyme variation in fishes. Echoing a sentiment popular at the time, I concluded that regulatory rather than structural genes should be studied next because changes in gene regulation were perhaps at the heart of adaptive evolution. I queried the audience for suggestions on how I might examine regulatory genes, and one responder asked whether I had considered using restriction enzymes to assay repetitive nuclear DNA

sequences, which at the time were viewed as prime candidates as regulatory modulators (Britten & Davidson 1969, 1971). I had never heard of restriction enzymes! However, the idea was intriguing so I soon approached several faculties at the University in an attempt to identify a collaborating laboratory where I might learn restriction digestion techniques. To my chagrin, the inquiries met with cool responses, except one: Dr Robert Lansman welcomed me to his laboratory, but noted with apology that he had limited experience with nuclear DNA and instead conducted research on the biochemistry and cellular biology of mitochondrial DNA. I barely had heard of mitochondrial DNA! However, left with few options, I accepted Bob's offer in order to gain familiarity with DNA level assays.

Before long, we were generating agarose gels with mtDNA restriction profiles, initially from small mammals. Although I was still viewing the effort mainly as a training exercise, intriguing questions began to emerge. Why did each individual display only a few mtDNA bands on a gel, rather than a smear of fragments from the billions of mtDNA molecules that must be included in an assay? (It must be because each specimen had a specifiable mtDNA genotype with respect to the restriction sites assayed.) Why did different mice within local populations often display distinct RFLP patterns, such that observed mtDNA variation primarily was distributed among rather than within individuals? (With hindsight, it must be because mtDNA mutations arise




Fig. 3 Breakdown of the phylogeographic articles from Fig. 1 according to the molecule or assay procedure employed.




Fig. 4 Examples of different 'allelic pathways' (Avice & Wollenberg 1997) within one-and-the-same hypothetical organismal pedigree. Upper left: the pedigree completely specified, with males (M) indicated by closed squares and females (F) by open circles. Two lines connect each offspring to its biological parents and thereby also identify mating partners. The horizontal plane can be interpreted as a linear spatial axis, in which case the lines provide an indication of offspring dispersal distances within each of the three or more extant geographical populations pictured. (For many real species, the spatial dimension is two-dimensional rather than linear.) Other boxes: five different gender-defined transmission pathways available to nuclear alleles through this pedigree. In each box, bold arrows highlight the coalescent tree for the transmission pathway specified. For example, the central panel of the top row highlights the coalescent tree for the matrilineal pathways ($F \rightarrow F \rightarrow F \rightarrow F \rightarrow \dots F$, e.g. for mtDNA) tracing back in time from extant individuals, and the upper right panel shows the analogous coalescent tree for the patrilineal pathways ($M \rightarrow M \rightarrow M \rightarrow M \rightarrow \dots M$, e.g. for the Y chromosome). Bottom row: examples of three more coalescent trees among more than 4×10^6 definable trees (as gender-defined transmission pathways) for this 21-generation pedigree.

frequently, and sometimes precipitate within a small number of animal generations a genotypic turnover in the population of mtDNAs in a germ-cell lineage from which the assayed soma were derived.) Why did mtDNA genotypes in organismal populations appear connectable to one another in phylogenetically intelligible ways? (Because intermolecular recombination must be rare or nonexistent in these maternally inherited molecules, such that the matrilineal histories of mutation events were recorded in extant mtDNA genotypes.) Why did members of sexually reproducing species usually group together by mtDNA genotypes when the evolutionary connecting agents of mating and genetic recombination

seemed not to apply to these asexually transmitted genomes? (Because, as we now know, coalescent processes ensure phylogenetic links among genotypes via vertical pathways of ancestry even in the absence of inter-lineage genetic exchange mediated by mating events.) What ramifications might stem from the heretical practice made possible by mtDNA of viewing haplotypes as clones and individual animals as OTUs (operational taxonomic units), in population genetic analyses? (The list of responses is now long.)

In general, many unorthodox perspectives on evolution eventually were to emerge from studies of mtDNA (reviewed in Avice 1991), but years would pass before

relatively clear answers to some of the questions listed above and others similar to them were to be forthcoming. The lag time reflected in part the difficulty experienced by many researchers (certainly by me) in reorienting thought away from the traditional Mendelian perspectives that applied so well, for example, to allozyme systems on which many of us had been trained.

My collaboration with the Lansman laboratory went well, and our first paper on mtDNA variation in a natural population soon appeared (Avise *et al.* 1979a), followed shortly thereafter by the first large-scale phylogeographic survey of any species based on mtDNA lineages (Avise *et al.* 1979b). The technical stage for these efforts had been set in the early 1970s through prior mtDNA research on several fronts. For example, Brown & Vinograd (1974) and Upholt & Dawid (1977) had demonstrated the feasibility of generating restriction enzyme cleavage maps for animal mtDNAs; Dawid & Blackler (1972), and Hutchinson *et al.* (1974) among others had documented predominant maternal inheritance for mtDNA in higher animals; and Upholt (1977) had developed a statistical procedure for estimating sequence divergence among mtDNA genotypes from comparisons of restriction digests. Furthermore, in the same year that our first phylogeographic works appeared in print, Brown *et al.* (1979) published an extremely influential article highlighting the unexpected fast pace of mtDNA sequence evolution as gauged by interspecies comparisons of higher primates.

In the late 1970s, excitement generated by the new mtDNA discoveries ran high. I remember pondering the many research possibilities, of which two of anecdotal interest can be mentioned. Early on, it occurred to me that mtDNA might be a wonderful tool for analysing the evolution of parthenogenetic vertebrates, for at least two reasons. First, all such unisexual biotypes were thought to have arisen through hybridization between sexual species, such that by utilizing mtDNA data it should be possible to identify the maternal parent taxon in each case. Second, because parthenogenetic taxa reproduce asexually, the history of maternal lineages within them should in principle be one-and-the-same as the entire organismal phylogeny (unlike the case for a sexual species). I remember reasoning that it would be safe to shelve these ideas for the moment in the belief that many years would elapse before any molecular biologists might dream of this 'obscure' biological application for mtDNA. I could not have been more wrong. One of the first mtDNA analyses of natural populations dealt with precisely these evolutionary issues in parthenogenetic lizards (Brown & Wright 1975, 1979; see below)! Eventually, my laboratory did examine evolutionary processes in gynogenetic and hybridogenetic fish complexes using mtDNA (reviewed in Avise *et al.* 1992), but only well after Wes Brown, Craig Moritz (Brown's postdoctoral researcher at

the time), and their associates had produced an important series of mtDNA papers on the origins and evolution of parthenogenetic reptiles and other unisexual vertebrates (e.g. Densmore *et al.* 1989; Echelle *et al.* 1989; Moritz 1991).

It also seemed evident to Bob Lansman and myself that mtDNA analyses of human populations would be of great interest. However, we elected not to pursue this topic. Personally, I was wary of the inevitable social and political fallout from whatever findings might be uncovered about the nature of genetic differences between human skin colour races, or between humans and great apes; and, in any event, it seemed likely that the necessary research would be accomplished by someone. Here, my crystal ball proved truer. An influential study on human mtDNA evolution soon appeared (Brown 1980), followed by a number of more extensive but also controversy generating mtDNA analyses of higher primate phylogeny (e.g. Ferris *et al.* 1981; Brown *et al.* 1982) and human geographical variation (notably by Cann *et al.* 1987; reviews in Nei & Roychoudhury 1993; Cavalli-Sforza *et al.* 1994; Takahata 1995).

I should digress from this personal account for a moment to relate the history of Wes Brown's involvement with mtDNA, because this traces the other major root of evolutionary interest in the molecule. The story began in 1968 when Brown went to Caltech as a graduate student and was introduced to mtDNA in the laboratories of Giuseppe Attardi and Jerome Vinograd, where mtDNA transcription and physical chemistry, respectively, were being studied. In 1971, Brown went to an exhibition of Max Escher paintings at the Los Angeles County Museum, where he happened to meet John Wright, the curator of the herpetology department. Wright was probably the most knowledgeable person in the world on *Cnemidophorus* lizards, and Brown's chance meeting with him that day was to lead to their collaborative studies on the evolutionary origins of parthenogenetic taxa from a genealogical perspective. Brown gathered the mtDNA data at Caltech from 1971 to 1973 but, as mentioned above, the first papers did not appear until several years later. After a postdoctoral stint at the University of California at San Francisco, Brown moved across the Bay in 1978 to join the Allan Wilson group at Berkeley. There he restructured and equipped the laboratory for studies of animal mtDNA, and among other efforts initiated the important research mentioned above on human genealogical evolution.

Returning to the developing story at the University of Georgia, in those early years another important event for me personally stemmed from a casual conversation over lunch. I was explaining to my colleague Dr Michael Clegg our recent findings on modes of inheritance and patterns of geographical variation in mtDNA for small mammals, and he mentioned that the issues raised

seemed analogous to those for surname evolution in many human societies. This simple comment struck home, and helped greatly in my otherwise tortuous transition from Mendelian to phylogenetic thinking at the intraspecific level. The surname analogy does indeed hold well (Avice 1989b). Just as sons and daughters 'inherit' their father's nonrecombined surname (before recent rule changes in some families), so too do progeny normally receive nonrecombined mtDNA from their mothers. Furthermore, much the way that mutations sometimes arise in surnames (my own name was a 19th century misspelling of 'Avis'), point mutations occasionally arise and cumulatively differentiate related mtDNA genotypes. Thus, mtDNA molecules record matrilineal histories much as surnames record patrilineal, except that the matrilineal records extend much further back in time (surnames were invented *de novo* only within the past few centuries).

These insights were new to me, but not completely so to the field. Beginning much earlier in the century, statistical demographers had studied the dynamics of surname turnover in human populations (Lotka 1931) using models that now could be applied often with little modification to gene lineages such as those provided by mtDNA (Schaffer 1970). Such models stimulated my own and my students' efforts to examine the theoretical ties between population demography and phylogeographic patterns within (Chapman *et al.* 1982; Avice *et al.* 1984, 1988; Avice

1995) and among (Neigel & Avice 1986) populations and species, and to address these expectations in a series of empirical mtDNA studies on a wide variety of organisms in nature (reviewed in Avice 1994). 'Coalescent theory' is the term now applied to the formal mathematical and statistical properties of gene genealogies (Kingman 1982; Watterson 1984; Donnelly & Tavaré 1986; Hudson 1990), and results from this discipline are highly relevant to molecular phylogeographic interpretations.

Several other important developments in the history of phylogeography are summarized in Table 1. In addition to these signal events, throughout the 1980s and 1990s there has been a burgeoning growth in the application of both genealogical theory and molecular data to phylogeographic analyses. This has included extensions and refinements of coalescent theory for populations of varying demographies (Hudson 1990; Slatkin & Hudson 1991; Rogers & Harpending 1992; Nee *et al.* 1995; Eller & Harpending 1996; see also several articles in this issue of *Molecular Ecology*), improvements in statistical and cladistic procedures for extracting phylogeographic information from empirical data on gene genealogies (e.g. Slatkin 1989; Neigel *et al.* 1991; Templeton *et al.* 1995; Templeton & Georgiadis 1996; this issue), and a great plethora of empirical applications primarily involving mtDNA (Fig. 3). Of course, progress in several related areas, not the least of which are molecular and computer technologies, have contributed significantly to the general scientific climate

Table 1 Brief chronology of some of the important developments in the history of phylogeography*

Date	Development
1974	Brown & Vinograd demonstrate how to generate restriction site maps for animal mtDNAs
1975	Watterson describes some basic properties of gene genealogies, marking the beginnings of modern coalescent theory
	Brown & Wright introduce mtDNA analysis to the study of the origins and evolution of parthenogenetic taxa
1977	Upholt develops the first statistical method to estimate mtDNA sequence divergence from restriction digest data
1979	Brown, George & Wilson document rapid mtDNA evolution
	Avice, Lansman & colleagues present the first substantive reports of mtDNA phylogeographic variation in nature
1980	Brown provides an initial report on human mtDNA variation
1983	Tajima and also Hudson initiate statistical treatments of the distinction between a gene tree and a population tree
1986	Birmingham & Avice initiate comparative phylogeographic appraisals of mtDNA for multiple codistributed species
1987	Avice & colleagues coin the word 'phylogeography', define the field, and introduce several phylogeographic hypotheses
	Cann & colleagues in the Wilson laboratory describe extensive global variation in human mtDNA
1989	Slatkin & Maddison introduce a method for estimating interpopulation gene flow from the phylogenies of alleles
1990	Avice & Ball introduce principles of genealogical concordance as a component of phylogeographic assessment
1992	Avice summarizes the first extensive compilation, involving multiple species and genetic assays, of phylogeographic patterns for a regional fauna
1994	Moritz promotes the conceptual distinction between 'shallow' vs. 'deep' intraspecific phylogenies by introducing the terms 'management units' and 'evolutionarily significant units' (see also Ryder 1986; Riddle 1996).
1996	Volumes edited by Avice & Hamrick, and by Smith & Wayne, summarize the many roles for molecular phylogeographic analysis in conservation biology
1998	Interest in phylogeography continues to flower, as evidenced, for example, by this special issue of <i>Molecular Ecology</i>

*Particularly with regard to molecular and statistical sides of the field, and with due apologies to numerous other contributors whose works were important but due to space limitations cannot be included here.

that permitted the flowering of phylogeographic studies during the last two decades.

The future of phylogeography

What does the future hold for phylogeography? I suspect that the field is still in a rapid phase of growth (Fig. 1) and, as presaged by this special issue of *Molecular Ecology*, that many more empirical studies on diverse organisms can be anticipated. There will also be a further expansion of interest in the utility of coalescent theory as a formal conceptual thread for tying together more coherently the micro- and macroevolutionary disciplines of population genetics and phylogenetics.

More specifically, I see ample room for the expansion of phylogeography in three areas, each tied to a distinct aspect of 'genealogical concordance' (Avice 1996b). These three facets of concordance and their phylogeographic relevances are listed next, with brief descriptions of how I envision each as an exciting frontier for further research efforts.

Genealogical concordance, aspect 'i'

Concordance in significant genealogical partitions across multiple unlinked loci within a species.

Relevance: helps to establish that the phylogenetic partitions in gene trees register deep as opposed to shallow historical partitions in an organismal phylogeny.

As already noted, the great majority of empirical genetic research into phylogeography has involved mtDNA (Fig. 3), yet the matrilineal pathways of ancestry registered by this molecule represent only a minuscule fraction of the total historical record within a sexual organismal pedigree (Fig. 4). In principle, much of the remainder of that history should be ensconced in autosomal gene trees through which alleles have been transmitted via both genders. However, few attempts have been made to estimate nuclear gene genealogies in a phylogeographic context (Aquadro *et al.* 1991; Bernardi *et al.* 1993; Burton & Lee 1994; Palumbi & Baker 1994).

At least two complications, one technical and one biological, typically arise in attempts to recover genealogical information from nuclear genes. The technical difficulty is in isolating DNA haplotypes, one at a time, from diploid organisms at single-copy loci. Only with nuclear haplotypes cleanly separated can molecular assays such as DNA sequencing or restriction site mapping then be used to recover the phase (coupling vs. repulsion) of multiple DNA sequence variants, and thereby facilitate treatments of the alleles by phylogenetic procedures analogous to those conducted routinely for mtDNA haplotypes (which nature purifies). Avice (1994; p. 134) lists several experimental approaches for isolating haplotypes at particular nuclear

loci. To this list can be added two recent PCR-based approaches (Hillis *et al.* 1996) that involve physical separation of nuclear haplotypes either by 'DGGE' (denaturing gradient gel electrophoresis; Lessa 1993) or by 'SSCP' (single-strand conformational polymorphism; Ortí *et al.* 1997).

The second complication in the recovery of nuclear gene trees at the intraspecific level is biological, and necessitates the presence of genomic regions that accumulate mutations rapidly yet are nearly free of intragenic recombination over the ecological or evolutionary timescales of interest. It remains to be seen how common such gene regions are, and how readily they can be identified and studied, but this certainly is an open frontier for meaningful inquiry.

On the conceptual front, a 'multilocus coalescent theory' is needed – one that considers the means, variances, and frequency distributions of genealogical information across multiple unlinked gene trees within an organismal pedigree. The development of such a theory might begin with consideration of the composite genealogical properties expected for independent neutral loci in random mating populations with specified demographic histories. Useful phylogeographic extensions then might involve populations spatially structured in various ways, and also might consider epistatic or otherwise non-neutral genes, or those that display partial linkage. The broader challenge will be to integrate the multilocus coalescent theory with empirical data to be gathered from multiple nuclear (and mitochondrial) gene genealogies within and among spatially arrayed natural populations.

Genealogical concordance, aspect 'ii'

Concordance in the geographical positions of significant gene-tree partitions across multiple codistributed species.

Relevance: strongly implicates shared historical biogeographic factors in shaping the deeper intraspecific phylogenies, often on a regional scale.

Only a few empirical molecular studies have attempted comparative phylogeographic assessments within each of multiple codistributed species (Bermingham & Avice 1986; Avice 1992; da Silva & Patton 1993; Wenink *et al.* 1994; Turner *et al.* 1996; Zink 1996; Patton *et al.* 1997; see also several articles in this issue). Yet, such comparative assessments offer perhaps the greatest hope for significant advances in understanding how organismal behaviour, and the demographic and natural histories of populations, can influence intraspecific phylogeographic patterns. In this comparative light, findings of genealogical concordance will be of interest as reflections of shared community histories (Cracraft 1988). But findings of non-congruent genetic patterns will be valuable also because

they may illuminate historical differences among species in levels of gene flow, responses to geographical barriers or selective gradients, rates of molecular evolution, effective population sizes, or other such molecular, ecological, and demographic factors (Avice *et al.* 1987b; Bowen & Avice 1990; Lamb *et al.* 1992; Zink 1996).

Genealogical concordance, aspect 'iii'

Concordance of molecular gene-tree partitions with geographical boundaries between traditionally recognized biogeographic provinces.

Relevance: strongly implicates shared historical biogeographic factors as shapers of intraspecific organismal phylogenies and species distributional patterns.

Study of the third aspect of genealogical concordance inevitably will draw molecular phylogeography into closer contact with other disciplines such as ecology, historical geography, and phylogenetic biology writ large. Preliminary results with several faunas in the southeastern USA suggest that significant phylogeographic 'breaks' within species may tend to align geographically with traditionally recognized boundaries between biogeographic provinces as inferred from historical geological data, or from concentrations in the distributional limits of species (Avice 1996b). This suggests that historical factors influence not only species compositions in regional communities, but also the spatial distributions of genealogically distinct populations within species. Quite apart from the conceptual challenges motivated by such observations, the phylogeographic data themselves can be of great utility as summaries of the population genetic resources that conservation biology seeks to preserve (Avice & Hamrick 1996).

Phylogeography as a recognizable discipline grew from recent historical roots in molecular genetic analyses of mtDNA, and in mathematical studies of coalescent processes that seemed necessary to capitalize upon this new class of genealogical information within species. However, the full impact of phylogeographic thought remains to be realized in the broader biological and geographical sciences. Phylogeography has had an auspicious start. The greatest benefits and opportunities for the field will continue to arise, as they have in the past, from phylogeography's central, integrative position within the evolutionary and ecological sciences.

Acknowledgements

Work from the Avice laboratory has been supported by the NSF. I wish to thank Drs Wes Brown, Richard Hudson, Warren Ewens, James Crow for sharing with me their thoughts on matters of history.

References

- Aquadro CF, Weaver AL, Schaeffer SW, Anderson WW (1991) Molecular evolution of inversions in *Drosophila pseudoobscura*: the amylase gene region. *Proceedings of the National Academy of Sciences USA*, **88**, 305–309.
- Avice JC (1989a) Gene trees and organismal histories: a phylogenetic approach to population biology. *Evolution*, **43**, 1192–1208.
- Avice JC (1989b) Nature's family archives. *Natural History*, **3**, 24–27.
- Avice JC (1991) Ten unorthodox perspectives on evolution prompted by comparative population genetic findings on mitochondrial DNA. *Annual Review of Genetics*, **25**, 45–69.
- Avice JC (1992) Molecular population structure and the biogeographic history of a regional fauna: a case history with lessons for conservation biology. *Oikos*, **63**, 62–76.
- Avice JC (1994) *Molecular Markers, Natural History and Evolution*. Chapman & Hall, New York.
- Avice JC (1995) Mitochondrial DNA polymorphism and a connection between genetics and demography of relevance to conservation. *Conservation Biology*, **9**, 686–690.
- Avice JC (1996a) Space and time as axes in intraspecific phylogeography. In: *Past and Future Rapid Environmental Changes: The Spatial and Evolutionary Responses of Terrestrial Biota* (eds Huntley B, Cramer W, Morgan AV, Prentice HC, Allen JRM), pp. 381–388. Springer-Verlag, New York.
- Avice JC (1996b) Toward a regional conservation genetics perspective: phylogeography of faunas in the southeastern United States. In: *Conservation Genetics: Case Histories from Nature* (eds Avice JC, Hamrick JL), pp. 431–470. Chapman & Hall, New York.
- Avice JC, Ball RM Jr (1990) Principles of genealogical concordance in species concepts and biological taxonomy. *Oxford Surveys in Evolutionary Biology*, **7**, 45–67.
- Avice JC, Hamrick JL (1996) *Conservation Genetics: Case Histories from Nature*. Chapman & Hall, New York.
- Avice JC, Wollenberg K (1997) Phylogenetics and the origin of species. *Proceedings of the National Academy of Sciences USA*, **94**, 7748–7755.
- Avice JC, Lansman RA, Shade RO (1979a) The use of restriction endonucleases to measure mitochondrial DNA sequence relatedness in natural populations. I. Population structure and evolution in the genus *Peromyscus*. *Genetics*, **92**, 279–295.
- Avice JC, Giblin-Davidson C, Laerm J, Patton JC, Lansman RA (1979b) Mitochondrial DNA clones and matriarchal phylogeny within and among geographic populations of the pocket gopher, *Geomys pinetis*. *Proceedings of the National Academy of Sciences USA*, **76**, 6694–6698.
- Avice JC, Neigel JE, Arnold J (1984) Demographic influences on mitochondrial DNA lineage survivorship in animal populations. *Journal of Molecular Evolution*, **20**, 99–105.
- Avice JC, Arnold J, Ball RM Jr, Birmingham E, Lamb T, Neigel JE, Reeb CA, Saunders NC (1987a) Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. *Annual Review of Ecology and Systematics*, **18**, 489–522.
- Avice JC, Reeb CA, Saunders NC (1987b) Geographic population structure and species differences in mitochondrial DNA of mouthbrooding marine catfishes (Ariidae) and demersal spawning toadfishes (Batrachoididae). *Evolution*, **41**, 991–1002.

- Avise JC, Ball RM, Arnold J (1988) Current versus historical population sizes in vertebrate species with high gene flow: a comparison based on mitochondrial DNA lineages and inbreeding theory for neutral mutations. *Molecular Biology and Evolution*, **5**, 331–344.
- Avise JC, Quattro JM, Vrijenhoek RC (1992) Molecular clones within organismal clones: mitochondrial DNA phylogenies and the evolutionary histories of unisexual vertebrates. *Evolutionary Biology*, **26**, 225–246.
- Bermingham E, Avise JC (1986) Molecular zoogeography of freshwater fishes in the southeastern United States. *Genetics*, **113**, 939–965.
- Bernardi G, Sordino P, Powers DA (1993) Concordant mitochondrial and nuclear DNA phylogenies for populations of the teleost fish *Fundulus heteroclitus*. *Proceedings of the National Academy of Sciences, USA*, **90**, 9271–9274.
- Bowen BW, Avise JC (1990) Genetic structure of Atlantic and Gulf of Mexico populations of sea bass, menhaden, and sturgeon: the influence of zoogeographic factors and life-history patterns. *Marine Biology*, **107**, 371–381.
- Britten RJ, Davidson EH (1969) Gene regulation for higher cells: a theory. *Science*, **165**, 349–357.
- Britten RJ, Davidson EH (1971) Repetitive and non-repetitive DNA sequences and a speculation on the origins of evolutionary novelty. *Quarterly Review of Biology*, **46**, 111–138.
- Brown WM (1980) Polymorphism in mitochondrial DNA of humans as revealed by restriction endonuclease analysis. *Proceedings of the National Academy of Sciences, USA*, **77**, 3605–3609.
- Brown WM, Vinograd J (1974) Restriction endonuclease cleavage maps of animal mitochondrial DNAs. *Proceedings of the National Academy of Sciences, USA*, **71**, 4617–4621.
- Brown WM, Wright JW (1975) Mitochondrial DNA and the origin of parthenogenesis in whiptail lizards (*Cnemidophorus*). *Herpetological Review*, **6**, 70–71.
- Brown WM, Wright JW (1979) Mitochondrial DNA analyses and the origin and relative age of parthenogenetic lizards (genus *Cnemidophorus*). *Science*, **203**, 1247–1249.
- Brown WM, George Jr M, Wilson AC (1979) Rapid evolution of animal mitochondrial DNA. *Proceedings of the National Academy of Sciences, USA*, **76**, 1967–1971.
- Brown WM, Prager EM, Wang A, Wilson AC (1982) Mitochondrial DNA sequences of primates: tempo and mode of evolution. *Journal of Molecular Evolution*, **18**, 225–239.
- Burton RS, Lee B-N (1994) Nuclear and mitochondrial gene genealogies and allozyme polymorphism across a major phylogeographic break in the copepod *Tigriopus californicus*. *Proceedings of the National Academy of Sciences, USA*, **91**, 5197–5201.
- Cann RL, Stoneking M, Wilson AC (1987) Mitochondrial DNA and human evolution. *Nature*, **325**, 31–36.
- Cavalli-Sforza LL, Menozzi P, Piazza A (1994) *The History and Geography of Human Genes*. Princeton University Press, Princeton, New Jersey.
- Chapman RW, Stephens JC, Lansman RA, Avise JC (1982) Models of mitochondrial DNA transmission genetics and evolution in higher eukaryotes. *Genetical Research*, **40**, 41–57.
- Cracraft J (1988) Deep-history biogeography: retrieving the historical pattern of evolving continental biotas. *Systematic Zoology*, **37**, 221–236.
- Dawid IB, Blackler AW (1972) Maternal and cytoplasmic inheritance of mtDNA in *Xenopus*. *Developmental Biology*, **29**, 152–161.
- Densmore LD III, Moritz CC, Wright JW, Brown WM (1989) Mitochondrial DNA analyses and the origin and relative age of parthenogenetic lizards (genus *Cnemidophorus*). IV. Nine *sexlineatus*-group unisexuals. *Evolution*, **43**, 969–983.
- Donnelly P, Tavaré S (1986) The ages of alleles and a coalescent. *Advances in Applied Probability*, **18**, 1–19.
- Echelle AA, Dowling TE, Moritz CC, Brown WM (1989) Mitochondrial-DNA diversity and the origin of the *Menidia clarkhubbsi* complex of unisexual fishes (Atherinidae). *Evolution*, **43**, 984–993.
- Eller E, Harpending HC (1996) Simulations show that neither population expansion nor population stationarity in a West African population can be rejected. *Molecular Biology and Evolution*, **13**, 1155–1157.
- Ferris SD, Wilson AC, Brown WM (1981) Evolutionary tree for apes and humans based on cleavage maps of mitochondrial DNA. *Proceedings of the National Academy of Sciences, USA*, **78**, 2432–2436.
- Hey J (1994) Bridging phylogenetics and population genetics with gene tree models. In: *Molecular Ecology and Evolution: Approaches and Applications* (eds Schierwater B, Streit B, Wagner GP, DeSalle R), pp. 435–449. Birkhäuser Verlag, Basel, Switzerland.
- Hillis DM, Moritz C, Mable BK (eds) (1996) *Molecular Systematics*, 2nd edn. Sinauer, Sunderland, MA.
- Hudson RR (1983) Testing the constant-rate neutral allele model with protein sequence data. *Evolution*, **37**, 203–217.
- Hudson RR (1990) Gene genealogies and the coalescent process. *Oxford Surveys in Evolutionary Biology*, **7**, 1–44.
- Hutchinson III CA, Newbold JE, Potter SS, Edgell MH (1974) Maternal inheritance of mammalian mitochondrial DNA. *Nature*, **251**, 536–538.
- Kingman JFC (1982) The coalescent. *Stochastic Processes and their Applications*, **13**, 235–248.
- Lamb T, Jones TR, Avise JC (1992) Phylogeographic histories of representative herpetofauna of the southwestern US mitochondrial DNA variation in the desert iguana (*Dipsosaurus dorsalis*) and the chuckwalla (*Sauromalus obesus*). *Journal of Evolutionary Biology*, **5**, 465–480.
- Lessa EP (1993) Analysis of DNA sequence variation at population level by polymerase chain reaction and denaturing gradient gel electrophoresis. *Methods in Enzymology*, **224**, 419–428.
- Lotka AJ (1931) Population analysis – the extinction of families – I. *Journal of the Washington Academy of Sciences*, **21**, 377–380.
- Maddison W (1995) Phylogenetic histories within and among species. In: *Experimental and Molecular Approaches to Plant Biosystematics* (eds Hoch PC, Stephenson AG), pp. 273–287. Missouri Botanical Garden, St. Louis.
- Moritz CC (1991) The origin and evolution of parthenogenesis in *Heteronotia binoei* (Gekkonidae): evidence for recent and localized origins of widespread clones. *Genetics*, **129**, 211–219.
- Moritz CC (1994a) Applications of mitochondrial DNA analysis in conservation: a critical review. *Molecular Ecology*, **3**, 401–411.
- Moritz CC (1994b) Defining ‘evolutionarily significant units’ for conservation. *Trends in Ecology and Evolution*, **9**, 373–375.
- Nee S, Holmes EC, Harvey PH (1995) Inferring population history from molecular phylogenies. *Philosophical Transactions of the Royal Society of London*, **B349**, 25–31.
- Nei M, Roychoudhury AK (1993) Evolutionary relationships of human populations on a global scale. *Molecular Biology and Evolution*, **10**, 927–943.

- Neigel JE, Avise JC (1986) Phylogenetic relationships of mitochondrial DNA under various demographic models of speciation. In: *Evolutionary Processes and Theory* (eds Nevo E, Karlin S), pp. 515–534. Academic Press, New York.
- Neigel JE, Ball RM Jr, Avise JC (1991) Estimation of single generation migration distances from geographic variation in animal mitochondrial DNA. *Evolution*, **45**, 423–432.
- Ortí G, Hare MP, Avise JC (1997) Detection and isolation of nuclear haplotypes by PCR–SSCP. *Molecular Ecology*, **6**, 575–580.
- Palumbi SR, Baker CS (1994) Contrasting population structure from nuclear intron sequences and mtDNA of humpback whales. *Molecular Biology and Evolution*, **11**, 426–435.
- Patton JL, da Silva MNE, Lara MC, Mustrangi MA (1997) Diversity, differentiation, and the historical biogeography of non-volant small mammals of the neotropical forests. In: *Tropical Forest Remnants: Ecology, Management, and Conservation of Fragmented Communities* (eds Laurance WF, Bierregaard RO Jr), pp. 455–465. University of Chicago Press, Chicago, Illinois.
- Riddle BR (1996) The molecular phylogeographic bridge between deep and shallow history in continental biotas. *Trends in Ecology and Evolution*, **11**, 207–211.
- Rogers AR, Harpending H (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution*, **9**, 552–569.
- Ryder OA (1986) Species conservation and the dilemma of subspecies. *Trends in Ecology and Evolution*, **1**, 9–10.
- Schaffer H (1970) The fate of neutral mutants as a branching process. In: *Mathematical Topics in Population Genetics* (ed. Kojima K), pp. 317–336. Springer-Verlag, New York.
- da Silva MNE, Patton JL (1993) Amazonian phylogeography: mtDNA sequence variation in arboreal echimyid rodents (Caviomorpha). *Molecular Phylogenetics and Evolution*, **2**, 243–255.
- Slatkin M (1989) Detecting small amounts of gene flow from phylogenies of alleles. *Genetics*, **121**, 609–612.
- Slatkin M, Maddison WP (1989) A cladistic measure of gene flow inferred from the phylogenies of alleles. *Genetics*, **123**, 603–613.
- Slatkin M, Hudson RR (1991) Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics*, **129**, 555–562.
- Smith TB, Wayne RK (eds) (1996) *Molecular Genetic Approaches in Conservation*. Oxford University of Press, New York.
- Tajima F (1983) Evolutionary relationship of DNA sequences in finite populations. *Genetics*, **105**, 437–460.
- Takahata N (1995) A genetic perspective on the origin and history of humans. *Annual Review of Ecology and Systematics*, **26**, 343–372.
- Templeton AR, Georgiadis NJ (1996) A landscape approach to conservation genetics: conserving evolutionary processes in the African bovidae. In: *Conservation Genetics: Case Histories from Nature* (eds Avise JC, Hamrick JL), pp. 398–430. Chapman & Hall, New York.
- Templeton AR, Routman E, Phillips CA (1995) Separating population structure from population history: a cladistic analysis of the geographical distribution of mitochondrial DNA haplotypes in the tiger salamander, *Ambystoma tigrinum*. *Genetics*, **140**, 767–782.
- Turner TF, Trexler JC, Kuhn DN, Robison HW (1996) Life-history variation and comparative phylogeography of darters (Pisces: Percidae) from the North American central highlands. *Evolution*, **50**, 2023–2036.
- Upholt WB (1977) Estimation of DNA sequence divergence from comparison of restriction endonuclease digests. *Nucleic Acids Research*, **4**, 1257–1265.
- Upholt WB, Dawid IB (1977) Mapping of mitochondrial DNA of individual sheep and goats: rapid evolution in the D loop region. *Cell*, **11**, 571–583.
- Watterson GA (1975) On the number of segregating sites in genetical models without recombination. *Theoretical Population Biology*, **10**, 256–276.
- Watterson GA (1984) Lines of descent and the coalescent. *Theoretical Population Biology*, **26**, 77–92.
- Wenink PW, Baker AJ, Tilanus MGJ (1994) Mitochondrial control-region sequences in two shorebird species, the turnstone and dunlin, and their utility in population genetic studies. *Molecular Biology and Evolution*, **11**, 22–31.
- Zink RM (1996) Comparative phylogeography in North American birds. *Evolution*, **50**, 308–317.

Work in the Avise laboratory, conducted by students and post-doctoral researchers, involves the application of molecular genetic markers to questions in natural history, ecology, and evolution. Avise himself spends most of his time dreaming up verbage to make it all seem more important.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.